

Breath-holding and its breakpoint

M. J. Parkes

School of Sport & Exercise Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK

This article reviews the basic properties of breath-holding in humans and the possible causes of the breath at breakpoint. The simplest objective measure of breath-holding is its duration, but even this is highly variable. Breath-holding is a voluntary act, but normal subjects appear unable to breath-hold to unconsciousness. A powerful involuntary mechanism normally overrides voluntary breath-holding and causes the breath that defines the breakpoint. The occurrence of the breakpoint breath does not appear to be caused solely by a mechanism involving lung or chest shrinkage, partial pressures of blood gases or the carotid arterial chemoreceptors. This is despite the well-known properties of breath-hold duration being prolonged by large lung inflations, hyperoxia and hypocapnia and being shortened by the converse manoeuvres and by increased metabolic rate.

Breath-holding has, however, two much less well-known but important properties. First, the central respiratory rhythm appears to continue throughout breath-holding. Humans cannot therefore stop their central respiratory rhythm voluntarily. Instead, they merely suppress expression of their central respiratory rhythm and voluntarily ‘hold’ the chest at a chosen volume, possibly assisted by some tonic diaphragm activity. Second, breath-hold duration is prolonged by bilateral paralysis of the phrenic or vagus nerves. Possibly the contribution to the breakpoint from stimulation of diaphragm muscle chemoreceptors is greater than has previously been considered. At present there is no simple explanation for the breakpoint that encompasses all these properties.

(Received 00 Month 2005; accepted after revision 31 October 2005; first published online 4 November 2005)

Corresponding author M. J. Parkes: School of Sport & Exercise Sciences, University of Birmingham, Edgbaston, Birmingham B15 2TT, UK. Email: m.j.parkes@bham.ac.uk

‘We can’t do, so we must think’¹

Introduction

The precise mechanisms explaining breath-holding and causing the breath at breakpoint are unknown. There are several useful reviews (Mithoefer, 1965; Godfrey & Campbell, 1968, 1969; Porter, 1970; Campbell & Guz, 1981; Lin, 1982; Nunn, 1987). Breath-holding is an unstable state with changes occurring in many interrelated variables. Although for clarity it is simplest to consider each important variable separately, the variables may well interact in ways that we cannot yet investigate by experiment.

This review will be restricted to experiments on humans while voluntarily breath-holding out of water. As yet there

is almost no comparable research on animals, because of the difficulty in persuading them to prolong the respiratory cycle even for more than ~5 s (Orem & Netick, 1986; Orem, 1989). Breath-hold diving (Lin, 1982) will not be considered, because face immersion evokes a diving reflex with different properties (Lin, 1982; Sterba & Lundgren, 1985; Butler & Woakes, 1987). Neither ‘involuntary’ breath-holding (e.g. central apnoea or suffocation) nor the sensation of breathlessness/air hunger/dyspnoea (Banzett *et al.* 1990; Ward *et al.* 2001) are considered because their precise relationships to the breakpoint are unclear.

There are many gaps in our understanding of breath-holding that raise difficulties in explaining the breakpoint.

The first difficulty is in how to best quantify breath-holding when the underlying control mechanisms are not known. Should measurement be related to potential sensory stimuli (of proprio- or chemoreceptors?), or to some subjective discomfort scale (which still begs the question of what stimulus causes the discomfort)? Without

¹Anonymous? paraphrased, *attr.* Lord Ernest Rutherford by Jones (1962).

a definitive answer, breath-hold duration (the time from the start of the breath-hold to the breakpoint breath) is easiest to quantify objectively.

The second difficulty is that even under similar experimental conditions, breath-holds produce discomfort which is not equally tolerated by all subjects. Hence breath-hold duration is variable, between studies, between subjects and within subjects (Fig. 1). Even within the same subject breath-hold duration can be increased by 13–19% with distractions [either by motor tasks (Fig. 2) or by mental arithmetic (Alpher *et al.* 1986)] or by 37% with successive trials (Fig. 3). This occurs even if inflation volume is the same for every breath and atelectasis is prevented by preceding each breath with a maximum lung inflation (Heath & Irwin, 1968). Experimental data on breath-hold duration should therefore be cited with some measure of variability (here, \pm s.e.m) and the effect of successive trials, distraction and motivation should always be considered. How can we ever be sure that all subjects are trying equally hard?




Figure 1. Variability in mean breath-hold duration between subjects, between studies and within subjects, or mean time to impending unconsciousness while breathing N₂

a, mean and range of breath-hold durations of 318 subjects in air at maximum inspiration (Schneider, 1930), used with permission of the American Physiological Society. *b*, mean and range of time breathing N₂ from eupnoea until impending unconsciousness of 45 subjects (Schneider, 1930), used with permission of the American Physiological Society. *c*, mean \pm 2s.d. breath-hold duration of 5 subjects in air at maximum inflation, used with permission of Lin *et al.* (1974) and the American Physiological Society. *d*, mean and range breath-hold duration of 2 subjects at functional residual capacity in 63% oxygen (Campbell *et al.* 1967), reproduced with permission from *Clinical Science* 1967, **32**, 425–432; © the Biochemical Society and the Medical Research Society.

The third difficulty, causing considerable confusion when comparing different studies, is that breath-hold duration depends on the experimental conditions (e.g. starting lung volume and inspired gas composition). These too must be cited.

Fourth, since many of the important studies used remarkably few subjects (not all of whom were naïve about physiology), or are unconfirmed or may be unrepeatable, it can be difficult to decide which are the most representative experiments.

Fifth, it is still not clear what happens to the respiratory musculature during breath-holding. Expiration in eupnoea is essentially a passive recoil process. Breath-holding is distinct from this because at large inflation volumes there may be some contribution from the voluntary muscles to hold the chest open at a chosen volume against this recoil. Holding cannot be explained simply by closure of the glottis and airway, because it is easy to continue the breath-hold with these structures open (Godfrey *et al.* 1969). The precise activity of the diaphragm, intercostal and accessory muscles during breath-holding has not been definitively established, but the diaphragm may contribute as the ‘holding’ muscle (see section entitled Paralysis of the diaphragm).

Sixth, although the simplest clue to the breakpoint mechanism should emerge from identifying any manoeuvre enabling breath-holding to unconsciousness, scientific reports of breath-holding to unconsciousness are rare and inconsistent, despite popular mythology. Schneider (1930) stated that ‘it is practically impossible for a man at sea level to voluntarily hold his breath until he




Figure 2. Percentage prolongation of mean breath-hold duration by various distractions

Prolongation by Valsalva, Mueller or ball squeezing manoeuvres in air at end expiratory volume (eupnoea), \pm s.e.m. in 6 subjects used with the permission of Bartlett (1977) and the American Physiological Society.

becomes unconscious, and subsequent scientific literature supports this in adults. [Anecdotal descriptions of losing consciousness describe subjects breath-holding at low barometric pressures, with low oxygen mixtures or with severe voluntary hyperventilation (Hill & Flacke, 1908; Schneider, 1930; Paulev, 1969). It may also be possible to cause unconsciousness by performing a Valsalva manoeuvre during breath-holding, although not reported by Bartlett (1977).] Even after the longest breath-holds from hyperoxia and hypocapnia, adult subjects break in an apparently involuntary manner and before any obvious impairment of cognitive function (Cooper *et al.* 2003), nor is any intellectual impairment obvious immediately afterwards. It follows from this that it is easiest to view the breath-hold as voluntary and the net stimulus that causes the breakpoint breath (generally an expiration followed by an inspiration) as usually involuntary, i.e. a powerful involuntary mechanism usually overrides the voluntary act of breath-holding. The breakpoint breath may be involuntary even when choosing to stop the breath-hold quite early, or when the breakpoint breath is only an attempt against a still closed airway. Viewing these in this way avoids a number of semantic difficulties, despite our lack of understanding of two crucial points: (a) the extent to which any breath is voluntary or involuntary (Shea, 1996); and (b) where the voluntary command to breath-hold intervenes in the pathway between the central respiratory rhythm generated in the brainstem (Feldman, 1986) and the spinal motoneurons of the respiratory muscles (Mitchell & Berger, 1975).




Figure 3. Prolongation of mean breath-hold duration by successive trials

Six subjects in 6 successive trials in air at end expiratory volume (eupnoea), \pm s.e.m used with the permission of Bartlett (1977) and the American Physiological Society.

Although the breakpoint breath is usually involuntary, some individuals claim to suppress it successfully. Even if they can, it would not appear to be a useful skill to acquire, since suppressing the breakpoint breath must ultimately lead to unconsciousness.

In the absence of manoeuvres that enable consistent breath-holding to unconsciousness, the next useful approach is to identify manoeuvres that prolong breath-hold duration. Since the cited breath-hold durations are so variable, perhaps a sensible guide is to consider manoeuvres important only if they almost double mean breath-hold duration.

Chest volume shrinkage and metabolic rate

The general effects on breath-hold duration of increasing lung inflation, inspired gas composition and metabolic rate are well known.

Breath-hold duration is increased by increasing lung inflation (Fig. 4*a*). It might be expected that lung volume stays constant throughout breath-holding when the extraction of O₂ from the alveoli is counteracted by equal production of CO₂ (i.e. presuming a respiratory exchange ratio of 1). In fact, as first shown by a decrease in buoyancy during breath-holding, the lungs gradually contract by 200–500 ml min⁻¹ during breath-holding (Stevens *et al.*




Figure 4. Effects of lung volume or exercise and lack of effect of pulmonary denervation or spinal anaesthesia on mean breath-hold duration

The lines indicate points to be compared and the error bars represent s.e.m. *a*, 10 normal subjects in air at total lung capacity (TLC) versus functional residual capacity (FRC), reprinted with permission from Flume *et al.* (1996) and from Elsevier. *b*, 5 control subjects versus 5 heart lung transplant patients in air at end expiratory volume, used with permission from Harty *et al.* (1996) and Blackwell Publishing. *c*, 4 subjects pre- versus post-T1 epidural anaesthesia in air at FRC, reproduced with permission from Eisele *et al.* (1968) and Noble *et al.* (1970), the Novartis Foundation and *Clinical Science* **35**, 23–33, © the Biochemical Society and the Medical Research Society. *d*, 5 subjects in air at maximum inflation, at rest versus during bicycle ergometry at 27 W, used with permission of Lin *et al.* (1974) and the American Physiological Society.

1946; Hong *et al.* 1971). This is because failure to remove CO₂ from the alveoli abolishes the partial pressure gradient that drives CO₂ from blood into alveolar gas, hence the extracted O₂ is not replaced by an equal volume of CO₂.

Since breath-hold duration depends on initial lung volume, which gradually decreases, one possible mechanism for the breakpoint might be the attainment of some minimum lung (or chest) volume causing sufficient sensory feedback to initiate a breath. (The precise afferents have never been specified since shrinkage classically unloads pulmonary stretch receptors.) Three types of experiment suggest that such a mechanism is not likely.

First, the breakpoint cannot be a simple function of lung shrinkage, because Godfrey *et al.* (1969) found that breath-hold duration was not shortened either if three subjects slowly exhaled during breath-holds or if the rate of lung shrinkage was increased by breath-holding at low ambient pressures.

Second, such a mechanism predicts that removing pulmonary afferents ought to influence breath-hold duration, if not to prolong it indefinitely. Neither is the case. Harty *et al.* (1996) showed that breath-hold duration was no different from control subjects in patients with pulmonary branches of the vagus nerve cut bilaterally following heart and lung transplantation (Fig. 4*b*), and Flume *et al.* (1996) found similar results.

Third, while pulmonary denervation alone does not eliminate all possible chest afferents, most of the remaining afferents ought to be eliminated by spinal anaesthesia. Eisele *et al.* (1968; Fig. 4*c*) found no effect on breath-hold duration of T1 epidural anaesthesia. This is the best evidence that afferents from the intercostal and from most of the accessory muscles of inspiration do not normally make an important contribution to breath-hold duration (unlike diaphragm afferents; see section entitled Paralysis of the diaphragm).

Increasing metabolic rate also shortens breath-hold duration (Rodbard, 1947; Cummings, 1962; Lin *et al.* 1974; Ward *et al.* 2001). Figure 4*d* shows that exercise (bicycle ergometry to at least double metabolic rate) more than halved breath-hold duration. Presumably, decreasing metabolic rate (e.g. by cooling or curarization) should prolong breath-hold duration.

Oxygen and carbon dioxide

During breath-holding, the arterial or end tidal partial pressure of oxygen P_{a/etO_2} falls below its normal level of ~100 mmHg and that of carbon dioxide $P_{a/etCO_2}$ rises above its normal level of ~40 mmHg. At breakpoint from maximum inflation in air, the P_{etO_2} is typically 62 ± 4 mmHg and the P_{etCO_2} is typically 54 ± 2 mmHg ($n = 5$; Lin *et al.* 1974), and the longer the breath-hold the more they change. It is remarkable that adults normally cannot breath-hold consistently to unconsciousness, even

under laboratory supervision. Nunn (1987) estimates that consciousness in normal adults is lost at P_{aO_2} levels below ~27 mmHg and P_{aCO_2} levels between 90 and 120 mmHg. Breakpoint levels close to these have been reported, e.g. P_{etO_2} levels as low as 24 mmHg, P_{etCO_2} levels as high as 91 mmHg and breath-hold durations of 14 min or more (Schneider, 1930; Ferris *et al.* 1946; Klocke & Rahn, 1959). For comparison, Schneider (1924, 1930) extraordinarily describes surreptitiously switching subjects' breathing to inspire from a spirometer of N₂ (and to exhale to room air) and measuring (Fig. 1*b*) the range of breathing times to impending unconsciousness (cyanosis, mask-like facial expression, pupil dilation, eye convergence, falling systolic pressure; Schneider & Truesdell, 1923). This range is similar to his range of breath-hold durations (Fig. 1*a*), yet such symptoms are not characteristic of the breakpoint of breath-holding.

One obvious hypothesis to explain the breakpoint is that once P_{aO_2} falls below or P_{aCO_2} rises above a certain threshold partial pressure, or rate of change of partial pressure reaches a threshold, then chemoreceptor stimulation causes an involuntary breath. The presumption has always been that these would be carotid chemoreceptors [aortic chemoreceptors have no demonstrable effect on breathing in humans (Lugliani *et al.* 1971; Wasserman *et al.* 1975)]. As the following paragraphs show, this 'arterial chemoreceptor hypothesis' is supported by the pronounced effects on breath-hold duration of altering the composition of the inspired gas. It is, however, confounded by the lack of a consistent pattern of arterial gas pressures at breakpoint, by denervation of carotid chemoreceptors failing to prolong breath-holds until unconsciousness and by the ability to breath-hold repeatedly after inspiring asphyxiating gas mixtures.

Breath-hold duration is almost doubled by breath-holding with hyperoxic gas mixtures (Fig. 5*c*), or by preceding breath-holding by voluntary or mechanical hyperventilation to lower P_{aCO_2} levels (Fig. 5*a*). Incidentally, preoxygenation has many practical advantages in studying breath-holding. Not only does it prolong duration, it also results in heart rate barely changing throughout the breath-hold (Gross *et al.* 1976) and in breakpoints that do not occur at P_{CO_2} levels low enough to threaten the brain. [Strictly, there is a risk of atelectasis with breath-holds when the lungs contain 100% O₂ (Campbell *et al.* 1967), so some dilution with nitrogen is preferable.]

Alternatively, breath-hold duration is almost halved by breath-holding from hypoxia (Fig. 5*c*), or from hypercapnia, e.g. raising the inspired P_{CO_2} to 65 mmHg (Godfrey & Campbell, 1969; Kelman & Wann, 1971).

The arterial chemoreceptor hypothesis, however, is not supported by the known blood gas pressures at breakpoint. Thus, preoxygenation does not prolong breath-hold duration until mean P_{aO_2} falls to *ca.* 62 mmHg.

Instead, the breakpoint occurs while P_{etO_2} is still remarkably elevated, e.g. 553 ± 16 mmHg, $n = 5$ (Lin *et al.* 1974). Conversely, hypoxia does not shorten breath-hold duration until P_{aO_2} falls to 62 mmHg. Instead the breakpoint occurs at the even lower P_{aO_2} values of 24–43 mmHg (Ferris *et al.* 1946). Similarly, hypercapnia does not shorten breath-hold duration until P_{etCO_2} rises to 54 mmHg (Kelman & Wann, 1971) and P_{etCO_2} can reach 70 mmHg (Godfrey & Campbell, 1969). Furthermore, the breakpoint of breath-holds from hypocapnia occurs at P_{etCO_2} levels between 48 ± 3 (Cooper *et al.* 2003) and 71 ± 3 mmHg (Klocke & Rahn, 1959). Nor is the breakpoint at some unique combination of low P_{etO_2} and high P_{etCO_2} (Klocke & Rahn, 1959). Indeed, even after the longest possible breath-holds from hypocapnia with preoxygenation, blood gas levels at breakpoint are remarkably benign.

In humans, the carotid bodies provide the only known means of detecting arterial hypoxia (Lugliani *et al.* 1971; Wasserman *et al.* 1975) and of rapidly detecting arterial hypercapnia. The arterial chemoreceptor hypothesis is further opposed by the fact that a breakpoint still occurs following carotid chemodenervation (resection),

i.e. denervation does not prolong breath-holding until unconsciousness. Davidson and coworkers (Davidson *et al.* 1974; Gross *et al.* 1976) compared breath-hold duration at inspiratory capacity in five patients following bilateral carotid body resection with that of normal subjects (Fig. 5*b*). Mean breath-hold duration in 100% O_2 was almost no different and there were no functionally important differences at breakpoint in their mean P_{etO_2} levels [362 ± 20 versus 425 ± 12 mmHg (mean \pm s.e.m.) in controls], nor in mean P_{etCO_2} levels [59 ± 2 versus 56 ± 4 mmHg (mean \pm s.e.m.)]. There can, however, be some ambiguity in interpreting these breath-hold duration data, because they can also be used to show that carotid body denervation does produce a small increase in breath-hold duration, confirmed by (Honda *et al.* 1988). Figure 5*b* shows that mean breath-hold duration in denervated patients in 21% O_2 is 54% longer ($P < 0.05$) than in intact subjects and that in 12% O_2 it is 65% longer ($P < 0.05$). Nevertheless, if the carotid chemoreceptors are the only means of detecting hypoxia, what mechanism explains how hypoxia continues to shorten breath-hold duration in denervated patients? Possibly, this shortening still occurs because the important action of hypoxia is




Figure 5. Effects of oxygenation or hypocapnia and of carotid denervation on mean breath-hold duration

Lines indicate the appropriate comparisons and the error bars represent s.e.m. *a*, 7 subjects breath-holding at maximal inspiration during normocapnia (41 ± 1 mmHg) or hypocapnia (24 ± 1 mmHg) in 100% O_2 (Klocke & Rahn, 1959), used with permission of the American Physiological Society. *b*, 7 intact versus 5 denervated subjects in 100, 21 and 12% O_2 at inspiratory capacity, used with permission from Davidson *et al.* (1974) and Gross *et al.* (1976), the American Physiological Society and the *New England Journal of Medicine*. *c*, 23 normal subjects breath-holding in 100, 21 and 10% O_2 at maximum inflation (Engel *et al.* 1946), reproduced with permission from *Journal of Clinical Investigation*.

not on carotid chemoreceptors but is on diaphragm muscle chemoreceptors (Road, 1990; Jammes & Speck, 1995), whose stimulation may instead make an important contribution to the breakpoint (see section entitled Paralysis of the diaphragm).

Do central chemoreceptors mediate the breakpoint? Their role during breath-holding is still unclear. In as much as P_{aCO_2} reflects their level of stimulation during breath-holding, the lack of a consistent P_{aCO_2} level at breakpoint suggests not. Yet the facts that breath-hold duration in 5 patients with apparently no functional peripheral or central chemoreceptivity (congenital central hypoventilation syndrome- Shea *et al.*, 1993) is almost double that of 5 age and gender matched controls, and that 4/5 had to be told to break by the experimenters, suggests otherwise.

Repeatedly breath-holding following inspiration of asphyxiating gas

The most dramatic demonstration that breath-hold duration is not simply dependent on blood gas pressures comes from measuring the effect at breakpoint of breathing asphyxiating gas mixtures (i.e. mixtures whose inspiration lowers P_{aO_2} and raises P_{aCO_2} even further) on the ability to make successive breath-holds. This appears to be not widely known, yet was clearly described 51 years ago by Fowler (1954) and is alluded to much earlier (Hill & Flacke, 1908).

If there exists some threshold partial pressure(s) that invokes the involuntary termination of breath-holding, subjects should not be able to make a second breath-hold without first restoring blood gas pressures to normal. Fowler (1954), however, showed that at breakpoint (mean end-tidal P_{etCO_2} 47 mmHg, $n = 3$ and mean oxygen saturation (S_{aO_2} –3% of control values), allowing eight subjects eight breaths of an asphyxiating mixture (8% O_2 and 7.5% CO_2) enabled them immediately to perform another breath-hold for 20 s. At the breakpoint of the second breath-hold (mean P_{etCO_2} 51 mmHg, $n = 3$ and mean S_{aO_2} –10% of control values), another eight breaths of the asphyxiating gas enabled a further 20 s breath-hold (with gases at breakpoint being a mean P_{etCO_2} of 52 mmHg, $n = 3$ and mean S_{aO_2} of –12%). This was subsequently confirmed with 24 subjects (Flume *et al.* 1994).

Not only is the ability to undertake a second breath-hold essentially independent of blood gas levels, it is also independent of the volume or number of the intervening involuntary breath(s) (Godfrey & Campbell, 1969; Rigg *et al.* 1974; Flume *et al.* 1994, 1995). This ability also persists if performing an isovolume manoeuvre, or merely an inspiratory effort (–12 cmH₂O pressure) against a closed airway (Rigg *et al.* 1974), and even after bilateral lung transplantation in nine subjects (Flume *et al.* 1996). The explanation for this ability may be that stopping

the voluntary breath-hold confounds the involuntary breakpoint mechanism, so another breath-hold is always possible. Confounding might be achieved simply as a result of relaxing any tonic diaphragm activity (see section entitled Paralysis of the diaphragm).

The central respiratory rhythm and breath-holding

The two vital rhythms in all animals are the heart's rhythm and the central respiratory rhythm. Strictly the term 'central respiratory rhythm' is the prerogative of the neurophysiologist, who can record both brainstem respiratory neurone activity and its output in phrenic motoneurons (Feldman, 1986). At present such phrenic recordings are not possible in humans. Nevertheless this term is used here for clarity, even though the closest approximation to its measurement in humans can only be from recording autonomic outflow or skeletal muscle activity.

Humans have almost no voluntary control of their heart beating. For instance, they cannot voluntarily pace their heart rate to a metronome, ~ or voluntarily change stroke volume, ~ or voluntarily stop their heart. Humans, however, have much voluntary control of breathing. They can easily pace their breathing rate to a metronome between 1 and 30 breaths min^{-1} and can voluntarily control tidal volume. It has been tacitly presumed that, because breath-holding in humans is voluntary, humans can voluntarily stop their central respiratory rhythm. Evidence has been available since the 1960s, however, showing that this is not the case.

In 1963 Agostoni asked three subjects to swallow catheters so that their tips lay in the oesophagus near the diaphragm, enabling measurement of intrathoracic pressure and indirect measurement of diaphragmatic electromyogram (EMG) activity (Agostoni, 1963). Agostoni showed that during breath-holding no EMG activity was detectable initially. Rhythmic negative pressure fluctuations and simultaneous rhythmic EMG activity appeared towards the end of breath-holding (Fig. 6A) and their frequency was within the respiratory frequency range. The amplitude and frequency of these pressure waves increase towards the breakpoint (Lin *et al.* 1974; Whitelaw *et al.* 1981, 1987). Corresponding activity can even be seen in some subjects during breath-holding as rhythmic 'tracheal tugging' against a closed glottis. The simplest explanation is that these are caused by the central respiratory rhythm. This is because the rhythmic EMG and negative pressure waves occur simultaneously, because their frequency and amplitude are within the respiratory range and because they increase as CO_2 levels rise towards the end of the breath-hold. This CO_2 rise would stimulate the central respiratory rhythm.

The appearance of the respiratory rhythm before the breakpoint shows additionally that the breakpoint itself


is not caused by the sudden restarting of the respiratory rhythm. The failure to record such rhythmic EMG activity throughout breath-holding, however, raises a problem: how to distinguish between the central respiratory rhythm being voluntarily stopped at the start of breath-holding and reappearing towards the breakpoint, or never stopping but merely being undetectable at the start of breath-holding?

Since there is no voluntary control of heart rate, one solution is use respiratory sinus arrhythmia (the decrease in heart period that usually accompanies every

breath) to indicate whether the central respiratory rhythm persists during breath-holding. We have recently shown that respiratory sinus arrhythmia in humans is caused predominantly by the central respiratory rhythm, rather than being a secondary effect of rhythmic chest inflation or negative intrathoracic pressure and its mechanical sequelae (Daly, 1986). This is because the central respiratory rhythm and sinus arrhythmia remain during mechanical hyperventilation in normocapnia but are greatly reduced in hypocapnia (Cooper *et al.* 2004). [The phenomenon of post-hyperventilation breathing or

Figure 6. Diaphragm activity and respiratory sinus arrhythmia during breath-holding

A, appearance of rhythmic diaphragm EMG activity towards the end of breath-holding. Oesophageal pressure (upper trace) and diaphragm EMG activity (lower trace) in one 22-year-old subject during breath-holding at resting lung volume in air, used with the permission of Agostoni (1963) and the American Physiological Society. The start and end of the breath-hold are indicated by circled arrows. **B**, sinus arrhythmia in panel **A** from the start of breath-holding. Using a copy of the original record from panel **A** kindly provided by Professor Agostoni, I have measured the time of each R wave (to within 20 ms) in each ECG artefact to calculate instantaneous heart period as described by Cooper *et al.* (2004) and measured the time and size of each of the 12 oesophageal pressure waves. The start and end of the breath-hold are indicated by arrows. **C**, some of the sinus arrhythmia in panel **A** is respiratory in origin. I have sampled the instantaneous heart period and pressure waves from panel **B** every 2.5% of time between the start of each pressure wave and the next. I then averaged both over the 11 oesophageal pressure wave intervals during the breath-hold as described by Cooper *et al.* (2004). Values plotted are means \pm s.e.m. This shows that some of the sinus arrhythmia is respiratory in origin, i.e. some sinus arrhythmia shows a similar relationship to rhythmic diaphragm activity (heart period decreases during inspiration) to that seen during eupnoea (see Fig. 1C of Cooper *et al.* 2004).



after-discharge (Tawadrous & Eldridge, 1974) is not detectable in such experiments (Cooper *et al.* 2003, 2004).]

There has been considerable debate over whether sinus arrhythmia persists during breath-holding (Angelone & Coulter, 1965; Davies & Neilson, 1967*a,b*; Valentinuzzi & Geddes, 1974; Hirsch & Bishop, 1981; Daly, 1986; Fritsch *et al.* 1991; Eckberg & Sleight, 1992; Pawelczyk & Levine, 1995; Trzebski & Smietanowski, 1996; Passino *et al.* 1997; Piepoli *et al.* 1997; Trzebski *et al.* 2001; Javorka *et al.* 2001). Part of this debate arises because subjects rarely breath-hold in air for longer than 1 min and often for only 20–45 s. This is barely long enough for a few central respiratory cycles to occur, never mind resolving whether or not sinus arrhythmia also persists and reveals them. Prolonging breath-hold times to *ca.* 4 min using preoxygenation is, however, sufficient to test whether sinus arrhythmia is

present and whether sinus arrhythmia behaves as if the central respiratory rhythm continues from the start of breath-holding. We have shown in such prolonged breath-holds in 10 subjects that sinus arrhythmia does persist from the start of breath-holding (e.g. Fig. 7*E*) and possesses the CO₂ sensitivity characteristic of the central respiratory rhythm (Cooper *et al.* 2003). When preparing this review, Professor Agostoni kindly gave me a copy of his original data (Fig. 6*A*) showing diaphragm activity with a large ECG artefact during breath-holding. Figure 6*B* shows that there is sinus arrhythmia from the start of breath-holding in these data too, and Fig. 6*C* shows that some of this sinus arrhythmia is respiratory in origin. [The sinus arrhythmia (Fig. 6*B*) is not obviously different when these pressure/EMG waves first appear, nor is it generally believed that sinus arrhythmia is suddenly




Figure 7. Sinus arrhythmia in the subject with the longest breath-hold (min) in normocapnia with preoxygenation at maximum inflation (Cooper *et al.* 2003), used with the permission of the American Physiological Society

A, air flow in eupnoea immediately preceding breath-holding. B, distribution of the 13 breath intervals in a within the 30 usable bins describing frequency (0.03–0.5 Hz). The dashed line indicates his eupneic frequency range (bins 7–10). C–L, instantaneous heart period and its $\text{power}_{0.03-0.5\text{Hz}}$ density spectra in eupnoea (C and D), the first 2 min (E and F) and last 2 min (G and H) of breath-holds from normocapnia (6 min. duration) and from hypocapnia (6.7 min duration) (I–L) [‘hypocapnia’ corrects the typographical error in the original Fig. 1 of (Cooper *et al.* 2003)]. The horizontal scale in B, D, F, H, I and J indicates each of the 32 bins describing 0–0.5 Hz. [The arrow in J indicates the lack of evidence for any short-term potentiation or after-discharge (Tawadrous & Eldridge, 1974) following mechanical hyperventilation because there is no increase in $\text{power}_{0.03-0.5\text{Hz}}$ at the frequency (bin 18) of the preceding ~20 min of mechanical hyperventilation.]

different towards the end of breath-holding (Cooper *et al.* 2003). It is not clear whether the moment these waves first appear, sometimes named the physiological breakpoint (Lin, 1982) as distinct from the conventional breakpoint, has any particular significance.]

In the absence of any more direct measures of the central respiratory rhythm being available in humans, all this evidence indicates that the central respiratory rhythm is present throughout breath-holding.

This conclusion has several important implications.

First, it shows that humans cannot voluntarily stop, i.e. that they do not have absolute control of, their central respiratory rhythm. Instead they breath-hold merely by voluntarily 'holding' the chest and suppressing expression of their central respiratory rhythm. Where and how voluntary suppression occurs remain unclear. Is the location the brainstem but effectively bypassing the respiratory input to cardiac vagal preganglionic neurones, or is it more distal, e.g. in the spinal cord? (Nathan & Sears, 1960; Mitchell & Berger, 1975). Is voluntarily 'holding' assisted by some tonic voluntary contraction of the diaphragm? Does this voluntary input to phrenic motoneurones continue during each expiratory phase of the central respiratory rhythm and override any potential inhibition from a central respiratory drive potential (Sears, 1966)? Such mechanisms, if they do operate in this way, could so easily explain how breath-holding suppresses expression of the central respiratory rhythm, without stopping the rhythm itself.

Second, it has caused confusion previously to describe a breath-hold as inspiratory, because this implies that the central respiratory rhythm has stopped in its inspiratory phase, or for an expiratory breath-hold to have stopped in its expiratory phase. Better terminology would be an inflation or a deflation breath-hold. Or is it only the starting lung volume as held by the diaphragm that is important? Is the means of reaching it irrelevant?

Third, some studies use breath-holding to study physiological mechanisms with the presumption that the central respiratory rhythm has stopped. This presumption appears invalid. Strictly, such studies only consider mechanisms in the absence of rhythmic pulmonary inflation. It may be invalid even for breath-holds from hypocapnia, since the hypocapnia levels safely attainable in humans would not necessarily stop the central respiratory rhythm as measured in animals (Boden *et al.* 1998).

Fourth, might all chemoreceptor contributions to the breakpoint be mediated through the central respiratory rhythm?

Paralysis of the diaphragm

Between 1967 and 1969 Campbell paralysed voluntary musculature with *d*-tubocurarine in two atropinized

subjects (one had an oral airway inserted) and mechanically ventilated them with 63% O₂ via a facepiece (Campbell *et al.* 1966, 1967, 1969). Voluntary control of one arm was retained (using an arterial occlusion cuff to restrict the entry of curare) to enable the subject to signal when they wanted to 'breathe'. The ventilator was then switched off and the conscious subjects were asked to signal when they wanted ventilation restarted. Their mean 'breath-hold' durations (Fig. 1*d*) were prolonged at least threefold (Campbell *et al.* 1967), and after ~4 min the experimenters intervened. P_{etCO_2} levels at 'breakpoint' of up to 72 mmHg were reported (Campbell *et al.* 1969). When they could again talk, subjects reported no distressing symptoms of suffocation or of discomfort.

Interpretation of this remarkable experiment has been challenged in two ways. First, Campbell's results were not confirmed. Gandevia *et al.* (1993) performed a similar experiment (except that in all 3 subjects the trachea was intubated transnasally) and found that curare did not prolong 'breath-hold' duration (mean duration of control breath-holds 79 s, range 48–110 s *versus* curarized 'breath-holds' of 78 s, range 34–120 s) and reported severe dyspnoea at 'breakpoint'. In neither study, however, could the subjects be termed naïve to possible outcomes! Secondly, curare must prolong 'breath-hold' duration to a small extent because it reduces metabolic rate.

Both challenges may be addressed. First, while greatly respecting the enormous courage of Gandevia's subjects in consciously placing their lives in the experimenters' hands, they may have suffered additional discomfort from intubation and their P_{etCO_2} levels at 'breakpoint' [43 ± 3 *versus* 43 ± 3 mmHg (means \pm s.d.) without paralysis] were almost normal. [The equally remarkable study by Banzett *et al.* (1990), of the perception of air hunger when awake, curarized subjects were mechanically ventilated continuously and P_{etCO_2} was changed in 3–4 mmHg steps, is so unlike the brief and non-steady state of breath-holding that it is not directly comparable. Nevertheless the study by Banzett *et al.* (1990) only eliminates the role of diaphragm/respiratory muscle contraction in air hunger and is still consistent with a possible role of diaphragm chemoreceptors in air hunger].

Secondly, any debate between Campbell's and Gandevia's remarkable, probably irreconcilable and unrepeatable studies can be side-stepped by considering the studies of Noble *et al.* (1970, 1971). They injected local anaesthetic into the phrenic nerves bilaterally, therefore restricting the paralysis to the diaphragm, i.e. without such a life-threatening manoeuvre on conscious subjects and with less effect on metabolic rate. Noble and coworkers achieved similar approximate doubling (Fig. 8*a*) of breath-hold duration in three subjects (but still did not prolong breath-hold duration to unconsciousness). Moreover, 'in all three phrenic block subjects the sensation [of

breath-holding] was decreased or altered'. Similar results were found by Eisele *et al.* (1972).

If Campbell's, Noble's and Eisele's experiments are accepted, the obvious interpretation is that: (a) curarization or phrenic paralysis prolongs breath-hold time by preventing any diaphragm contraction and hence reduce generation of afferent feedback; and (b) it is afferent feedback from the diaphragm that normally causes the feeling of discomfort and may oppose the ability to breath-hold.

This interpretation however, raises several questions.

(1) What known afferent pathways can explain these effects? Diaphragm afferents travelling in the phrenic nerves are one obvious pathway. Although traditional respiratory neurophysiology (Feldman, 1986) does not normally consider that feedback from the diaphragm directly modulates the central respiratory rhythm, there is now substantial evidence that phrenic afferents may modulate diaphragm activity (Road, 1990; Jammes & Speck, 1995). Approximately 30% of the phrenic nerve in cats, dogs and rats carries afferents, i.e. 500–800 afferents per nerve (Ferguson, 1891; Landau *et al.* 1962; Langford &

Schmidt, 1983). Some of these are muscle proprioceptors (Corda *et al.* 1965; Balkowiec *et al.* 1995), but the majority appear to be type III or type IV afferents (Landau *et al.* 1962; Duron & Condamin, 1970; Langford & Schmidt, 1983). Some of these are active during spontaneous breathing, have receptive fields in the diaphragm and are stimulated by chemoreceptor stimulants (e.g. potassium, lactic acid, capsaicin or phenyl diguanide, or asphyxia or fatigue); in turn these stimuli, or electrical stimulation of phrenic afferents, may have reflex excitatory and/or inhibitory effects on the diaphragm (Graham *et al.* 1986; Jammes *et al.* 1986; Supinski *et al.* 1989, 1993; Balkowiec *et al.* 1995; Iscoe & Duffin, 1996). Little is known about the role of diaphragm afferents in humans, although there is indirect evidence that they may be important (Nathan & Sears, 1960; Green *et al.* 1978). Strictly, however, the experiments of Campbell, Noble and Eisele establish only the effects on breath-hold duration of motor paralysis of the diaphragm. Without selective blockade of phrenic afferents only, they do not establish that the afferent pathway is via the phrenic nerves.

(2) What does this afferent feedback indicate? Does it indicate proprioceptor activity (the presence of any tonic diaphragmatic contraction, or the absence of rhythmic diaphragm contractions), or merely diaphragm fatigue (i.e. diaphragm chemoreceptor activity; Fisher & White, 2004)? Without knowing the precise status of the respiratory muscles during breath-holding this is unclear. To explain how the chest is held inflated at large volumes during breath-holding, even with an open glottis and airway, it might be presumed that there is some tonic activation of the diaphragm during breath-holding. Yet in humans there is no direct evidence about what breath-holding does to phrenic motoneurone activity. EMG activity is usually recorded only indirectly from what is topologically the body surface during breath-holding (Agostoni, 1963), so it is hardly surprising that only rhythmic diaphragm EMG activity is detected and only towards the end of breath-holding (Fig. 6A). The possibility remains that voluntarily 'holding' involves some tonic and almost isometric diaphragm contraction ('almost' because of lung shrinkage throughout breath-holding). Such a mild contraction may not always be discernible as a diaphragm EMG signal detected on the body surface. Could such an unusual isometric contraction induce diaphragm fatigue sufficient to stimulate diaphragm muscle chemoreceptor afferents which contribute to the breakpoint. Even if the diaphragm does not contract tonically during breath-holding, some stimulation of diaphragm chemoreceptors is an intriguing alternative to the old arterial chemoreceptor hypothesis and may make more sense of many disparate observations. It could explain the effects of blood gas levels on breath-hold duration, if such muscle chemoreceptor activity




Figure 8. Prolongation of breath-hold duration mean \pm s.e.m by bilateral phrenic block or by bilateral vagus and glossopharyngeal nerve block

Prolongation of breath-hold duration by bilateral phrenic block (a), reproduced with permission from Noble *et al.* (1970, 1971), the Novartis Foundation and *Clinical Science* **41**, 275–283, © the Biochemical Society and the Medical Research Society, or bilateral vagus and glossopharyngeal nerve block (b), reproduced with permission from Noble *et al.* (1970) and the Novartis Foundation (and see also Guz *et al.* 1966; Guz, 1966). Note that breath-holds were at end expiration (FRC). They were from 100% O₂ for (a) (Noble *et al.* 1971) and apparently mostly from 100% O₂ for (b) (Guz *et al.* 1966; Guz, 1966; Noble *et al.* 1970). Squares and circles indicate individual subjects with the means \pm s.e.m. for each pre- and post-block condition indicated as horizontal bars.

is increased by arterial hypoxia and hypercapnia and decreased by hyperoxia and hypocapnia. It could also explain how stopping a voluntary breath-hold confounds the involuntary breakpoint mechanism, if stopping reduces stimulation of both muscle chemoreceptors (by restoring muscle blood flow) and proprioceptors.

(3) How is this feedback perceived? Normally humans have almost no sense of diaphragm *per se*, so precisely what is perceived from diaphragm muscle afferents during breath-holding is unclear. Even if the muscle afferent signal is only perceived vaguely as discomfort, the ability to tolerate such discomfort could explain the notorious variations in breath-hold duration (cf. the variation in holding times for isometric contractions in other voluntary muscles).

(4) Where are its principal integration sites? The facts that the central respiratory rhythm continues during breath-holding and that the breakpoint breath is usually involuntary suggest that the involuntary respiratory rhythm in the brainstem is the obvious principal site for integration. It is also possible, however, that this diaphragm feedback is important in opposing voluntary control of breathing and may act cortically or at a spinal level (Nathan & Sears, 1960). There is evidence for something similar to such feedback reaching consciousness in humans. This is the fact that when awake and mechanically hyperventilated in hypocapnia, volunteers almost never stop breathing (apnoeas ≤ 12 s) when the ventilator is switched off (Shea, 1996; Cooper *et al.* 2004). Yet when asleep, switching off in hypocapnia always produces apnoea ≥ 79 s (Henke *et al.* 1988; Datta *et al.* 1991).

Anaesthetic blockade of the vagus nerves

Guz, Noble and coworkers injected lignocaine bilaterally into the vagus (X) and glossopharyngeal (IX) nerves of three normal volunteers to block all afferent sensory traffic (Guz, 1966; Guz *et al.* 1966; Noble *et al.* 1970). They showed (Fig. 8*b*) that this alleviated the distress of breath-holding and, while not prolonging breath-hold duration indefinitely, still approximately trebled their mean breath-hold duration. (N.B. Fig. 8*b* breath-holds apparently were mostly from 100% O₂ and they appear short only because they were started from end expiration, whereas those in Fig. 5*b* were from inspiratory capacity.) Noble *et al.* (1970) made a further and ingenious connection:

‘The sensation of breath-holding was alleviated by vagal block, phrenic block, transection at the 3rd cervical segment and poliomyelitis. The sensation arises from frustrated contractions of the diaphragm stimulated by a reflex with its afferent limb in the vagus.’

This study has been neglected because such anaesthesia also blocks the afferents (Davidson *et al.* 1974) from the carotid (IX) and aortic (X) chemoreceptors and from the lungs (X). But we now know that there appears to be no aortic chemoreceptor contribution to breathing in humans, that the arterial chemoreceptor contribution to breath-hold duration is negligible in 100% O₂ (Fig. 5*b*), and that section of vagal afferents from the lungs has no effect (Fig. 4*b*) on breath-hold duration. The study by Noble and coworkers therefore deserves reconsideration. Their conclusion, ‘it thus appears that the build up of the stimulus to diaphragmatic contractions during breath-holding is dependent on vagal afferent information. We would welcome suggestions as to the precise nature of




Figure 9. One possible scheme for breath-holding and its breakpoint

this afferent mechanism' poses a challenge that no one appears to have taken up. Their experiment is particularly important because, since vagal blockade approximately trebled breath-hold duration while the phrenic afferent pathway was still intact, it implies that [non-pulmonary] vagus nerves may and phrenic nerves may *not* provide the principal afferent pathway from the diaphragm!

A unifying hypothesis for the breakpoint of breath-holding?

The simplest 'single variable' hypotheses based on either pressure levels of arterial blood gases or lung volume can be ruled out. Strictly, hypotheses based on various combinations of blood gases with lung volume (Mithoefer, 1965) cannot be excluded, but the complexity of multiple variable hypotheses makes them difficult to test scientifically.

Campbell and coworkers (Godfrey & Campbell, 1968, 1970; Rigg *et al.* 1974) consider the breakpoint in terms that can be simplified to a mismatch (inappropriateness) of return for effort (tension/length). These terms have still not been rigorously defined, but experiments since 1970 enable a further attempt (see Fig. 9).

Now that we know the central respiratory rhythm continues during breath-holding (and that the factors that increase it usually promote the breakpoint), defining these terms becomes more complex. The 'return' appears to involve muscle chemo- and proprioceptor afferent activity from the diaphragm. This may be perceived consciously only as the discomfort that opposes breath-holding. Nevertheless, since neither phrenic nor vagal blockade enables breath-holding to unconsciousness, are we still missing a crucial element of 'return'?

The 'effort' is to perform the unusual act of voluntarily 'holding' the chest (with the a contribution from diaphragm?). It might simply be a corticospinal input to phrenic motoneurons that bypasses the premotor respiratory neurones in the brainstem. In contrast, the breath that identifies the breakpoint is usually involuntary so presumably is not involved in this 'effort'.

We are, however, still no further in understanding what the term 'mismatch' means. Is the mismatch between the size of the effort *versus* return as the combined size of the diaphragm afferent signal with the central respiratory rhythm? Whatever it is, the smaller the mismatch the longer the breath-hold. Conversely, the bigger the mismatch the more likely is the breakpoint. And the breakpoint breath must cause the mismatch to disappear.

Treating mismatch in this way could explain a number of disparate properties of breath-holding, as follows.

First, mismatch is reduced by anything that reduces feedback from diaphragm afferents (e.g. curare, phrenic and possibly vagal block, arterial hyperoxia and hypocapnia) or that reduces the central respiratory rhythm

(e.g. arterial hyperoxia or hypocapnia, or increasing lung volume to increase the O₂ and CO₂ reservoir, or decreasing metabolic rate). These will prolong breath-hold time.

Second, mismatch is increased by anything that increases feedback from diaphragm afferents (e.g. any tonic diaphragm activity and possibly arterial hypoxia and hypercapnia) or that increases the central respiratory rhythm (e.g. arterial hypoxia or hypercapnia, or decreasing lung volume, or increased metabolic rate). These will shorten breath-hold time.

Third, because the breakpoint breath (releasing any tonic diaphragm contraction?) causes mismatch to disappear, another breath-hold is always possible.

Overall, considering a greater role for the diaphragm appears to provide a better explanation for the breakpoint, despite the lack of substantial supporting experimental evidence. This is why the paraphrase of Rutherford may be apt: 'We can't do, so we must think'. Perhaps recent developments in non-invasive imaging will improve our understanding of mismatch, effort and return.

References

- Agostoni E (1963). Diaphragm activity during breath holding: factors related to its onset. *J Appl Physiol* **18**, 30–36.
- Alpher VS, Nelson RB & Blanton RL (1986). Effects of cognitive and psychomotor tasks on breath-holding span. *J Appl Physiol* **1149**, 1152.
- Angelone A & Coulter NA (1965). Heart Rate Response to Held Lung volume. *J Appl Physiol* **20**, 464–468.
- Balkowiec A, Kukula K & Szulczyk P (1995). Functional classification of afferent phrenic nerve fibres and diaphragmatic receptors in cats. *J Physiol* **483**, 759–768.
- Banzett RB, Lansing RW, Brown R, Topulos GP, Yager D, Steele SM, Londono B, Loring SH, Reid MB, Adams L & Brown R (1990). 'Air hunger' from increased P_{CO₂} persists after complete neuromuscular block in humans. *Resp Physiol* **81**, 1–17.
- Bartlett D Jr (1977). Effects of Valsalva and Mueller maneuvers on breath-holding time. *J Appl Physiol* **42**, 717–721.
- Boden AG, Harris MC & Parkes MJ (1998). The apneic threshold for CO₂ in the anesthetized rat: fundamental properties under steady state conditions. *J Appl Physiol* **85**, 898–907.
- Butler PJ & Woakes AJ (1987). Heart rate in humans during underwater swimming with and without breath-hold. *Resp Physiol* **69**, 387–399.
- Campbell EJM, Freedman S, Clark TJ, Robson JG & Norman J (1966). Effect of curarisation on breath-holding time. *Lancet* **ii**, 207.
- Campbell EJM, Freedman S, Clark TJ, Robson JG & Norman J (1967). The effect of muscular paralysis induced by tubocurarine on the duration and sensation of breath-holding. *Clin Sci* **32**, 425–432.
- Campbell EJM, Godfrey S, Clarke JH, Freedman S & Norman J (1969). The effect of muscle paralysis induced by tubocurarine on the duration and cessation of breath holding during hypercapnia. *Clin Sci* **36**, 323–328.

- Campbell EJM & Guz A (1981). Breathlessness. In *Regulation of Breathing*, part II, ed. Hornbein TF, pp. 1181–1195. Marcel Dekker Inc., New York.
- Cooper HE, Clutton-Brock TH & Parkes MJ (2004). The contribution of the respiratory rhythm to sinus arrhythmia in normal unanesthetized subjects during mechanical hyperventilation with positive pressure. *Am J Physiol* **286**, H402–H411.
- Cooper HE, Parkes MJ & Clutton-Brock TH (2003). CO₂-dependent components of sinus arrhythmia from the start of breath-holding in humans. *Am J Physiol Heart Circ Physiol* **285**, H841–H848.
- Corda M, von Euler C & Lennerstrand G (1965). *J Physiol* **178**, 161–177. Proprioceptive innervation of the diaphragm.
- Cummings EG (1962). Breath holding at beginning of exercise. *J Appl Physiol* **17**, 221–224.
- Daly M de B (1986). Interactions between respiration and circulation. In *Handbook of Physiology*, section 3, *the Respiratory System*, vol. II, *The Control of Breathing*, ed. Fishman AP, pp. 529–594. American Physiological Society, New York.
- Datta AK, Shea SA, Horner RL & Guz A (1991). The influence of induced hypocapnia and sleep on the endogenous respiratory rhythm in humans. *J Physiol* **440**, 17–33.
- Davidson JT, Whipp BJ, Wasserman K, Koyal SN & Lugliani R (1974). Role of carotid bodies in breath holding. *New Eng J Med* **290**, 819–822.
- Davies CTM & Neilson JM (1967a). Disturbance of heart rhythm during recovery from exercise in man. *J Appl Physiol* **22**, 943–946.
- Davies CTM & Neilson JM (1967b). Sinus arrhythmia in man at rest. *J Appl Physiol* **22**, 947–955.
- Duron B & Condamin M (1970). Electron microscope study of the unmyelinated composition of the phrenic nerve of the cat. (In French). *Comptes Rendus des Seances la Societe Biologie de Ses Filiales* **164**, 577–583.
- Eckberg DL & Sleight P (1992). *Human Baroreflexes in Health and Disease*. Clarendon Press, Oxford.
- Eisele JH, Noble MIM, Katz J, Fung DL & Hickey RF (1972). Bilateral phrenic-nerve block in man: technical problems and respiratory effects. *Anesthesiology* **37**, 64–69.
- Eisele JH, Trenchard D, Burki N & Guz A (1968). The effect of chest wall block on respiratory sensation and control in man. *Clin Sci* **35**, 23–33.
- Engel GL, Ferris EB, Webb JP & Stevens CD (1946). Voluntary breath holding II. The relation of the maximum hold time to the oxygen tension of the inspired air. *J Clin Invest* **25**, 729–733.
- Feldman JL (1986). Neurophysiology of breathing in mammals. In *Handbook of Physiology*, section 1, *The Nervous System*, vol. 4, *Intrinsic Regulatory Systems of the Brain*, ed. Bloom FE, chapter 9, pp. 463–523. American Physiological Society, Washington, DC.
- Ferguson J (1891). The phrenic nerve. *Brain* **14**, 282–283.
- Ferris EB, Engel GL, Stevens CD & Webb A (1946). Voluntary breath-holding III. *J Clin Invest* **25**, 734–780.
- Fisher JP & White MJ (2004). Muscle afferent contributions to the cardiovascular response to isometric exercise. *Exp Physiol* **89**, 639–646.
- Flume PA, Eldridge FL, Edwards LJ & Houser LM (1994). The Fowler breathholding study revisited: continuous rating of respiratory sensation. *Resp Physiol* **95**, 53–66.
- Flume PA, Eldridge FL, Edwards LJ & Houser LM (1995). Relief of distress of breathholding: separate effects of expiration and inspiration. *Resp Physiol* **101**, 41–46.
- Flume PA, Eldridge FL, Edwards LJ & Mattison LE (1996). Relief of the ‘air hunger’ of breathholding. A role for pulmonary stretch receptors. *Resp Physiol* **103**, 221–232.
- Fowler WS (1954). Breaking point of breath holding. *J Appl Physiol* **6**, 539–545.
- Fritsch JM, Smith ML, Simmons DTF & Eckberg DL (1991). Differential baroreflex modulation of human vagal and sympathetic activity. *Am J Physiol* **260**, R635–R641.
- Gandevia SC, Killian KJ, McKenzie DK, Crawford M, Allen GM, Gorman RB & Hales JP (1993). Respiratory sensations, cardiovascular control, kinaesthesia and transcranial stimulation during paralysis in humans. *J Physiol* **470**, 85–107.
- Godfrey S & Campbell EJM (1968). The control of breath holding. *Resp Physiol* **5**, 385–400.
- Godfrey S & Campbell EJM (1969). Mechanical and chemical control of breath holding. *Q J Exp Physiol Cognate Med Sciences* **54**, 117–128.
- Godfrey S & Campbell EJM (1970). The role of afferent impulses from the lung and chest wall in respiratory control and sensation. In *Breathing; Hering–Breuer Centenary Symposium*, ed. Porter R, pp. 219–232. J. & A Churchill (Longmans Group), London.
- Godfrey S, Edwards RH & Warrell DA (1969). The influence of lung shrinkage on breath holding time. *Q J Exp Physiol Cognate Med Sciences* **54**, 129–140.
- Graham R, Jammes Y, Delpierre S, Grimaud C & Roussos C (1986). The effects of ischemia, lactic acid and hypertonic sodium chloride on phrenic afferent discharge during spontaneous diaphragmatic contraction. *Neurosci Lett* **67**, 257–262.
- Green M, Mead J & Sears TA (1978). Muscle activity during chest wall restriction and positive pressure breathing in man. *Resp Physiol* **35**, 283–300.
- Gross PM, Whipp BJ, Davidson JT, Koyal SN & Wasserman K (1976). Role of the carotid bodies in the heart rate response to breath holding in man. *J Appl Physiol* **41**, 336–340.
- Guz A (1966). Effects of blocking the vagus nerves in Man. In *Breathlessness*, ed. Howell JBM & Campbell EJM, pp. 65–71. Blackwells, Oxford.
- Guz A, Noble MIM, Widdicombe JG, Trenchard D, Mushin WW & Makey AR (1966). The role of vagal and glossopharyngeal afferent nerves in respiratory sensation, control of breathing and arterial pressure regulation in conscious man. *Clin Sci* **30**, 161–170.
- Harty HR, Mummery CJ, Adams L, Banzett RB, Wright IG, Banner NR, Yacoub MH & Guz A (1996). Ventilatory relief of the sensation of the urge to breathe in humans: are pulmonary receptors important? *J Physiol* **490**, 805–815.
- Heath JR & Irwin CJ (1968). An increase in breath-hold time appearing after breath-holding. *Resp Physiol* **4**, 73–77.

- Henke KG, Arias A, Skatrud JB & Dempsey JA (1988). Inhibition of inspiratory muscle activity during sleep. *Am Rev Resp Dis* **138**, 8–15.
- Hill L & Flacke JW (1908). The effect of excess carbon dioxide and of want of oxygen upon the respiration and circulation. *J Physiol* **37**, 77–111.
- Hirsch JA & Bishop B (1981). Respiratory sinus arrhythmia in humans: how breathing pattern modulates heart rate. *Am J Physiol* **241**, H620–H629.
- Honda Y, Hashizume I, Kimura H & Severinghaus JW (1988). Bilateral carotid body resection in man enhances hypoxic tachycardia. *Jap J Physiol* **38**, 917–928.
- Hong SK, Lin YC, Lally DA, Yim BJB, Kominami N, Hong PW & Moore TO (1971). Alveolar gas exchanges and cardiovascular functions during breath holding with air. *J Appl Physiol* **30**, 540–547.
- Iscoe S & Duffin J (1996). Effects of stimulation of phrenic afferents on cervical respiratory interneurons and phrenic motoneurons in cats. *J Physiol* **497**, 803–812.
- Jammes Y, Buchler B, Delpierre S, Rasidakis A, Grimaud C & Roussos C (1986). Phrenic afferents and their role in inspiratory control. *J Appl Physiol* **60**, 854–860.
- Jammes Y & Speck DF (1995). Respiratory control by diaphragmatic and respiratory muscle afferents. In *Regulation of Breathing*, 2nd edn, ed. Dempsey JA & Pack AI, chapter 12, pp. 543–582. Marcel Dekker, New York.
- Javorka M, Zila I, Javorka K & Calkovska A (2001). ‘Respiratory’ oscillations of cardiovascular parameters during voluntary apnea. *Resp Physiol* **126**, 251–254.
- Jones RV (1962). Science, technology and civilisation. *Bull Institute Physics* **12**, 97–103.
- Kelman GR & Wann KT (1971). Mechanical and chemical control of breath holding. *Q J Exp Physiol* **56**, 92–100.
- Klocke FJ & Rahn H (1959). Breath holding after breathing oxygen. *J Appl Physiol* **14**, 689–693.
- Landau BR, Akert K & Roberts TS (1962). Studies on the innervation of the diaphragm. *J Comparative Neurol* **119**, 1–10.
- Langford LA & Schmidt RF (1983). An electron microscopic analysis of the left phrenic nerve in the rat. *Anat Record* **205**, 207–213.
- Lin YC (1982). Breath-hold diving in terrestrial mammals. *Exercise Sport Sci Rev* **10**, 270–307.
- Lin YC, Lally DA, Moore TA & Hong SK (1974). Physiological and conventional breath-hold break points. *J Appl Physiol* **37**, 291–296.
- Lugliani R, Whipp BJ, Seard C & Wasserman K (1971). Effect of bilateral carotid body resection on ventilatory control at rest and during exercise in man. *New Eng J Med* **285**, 1105–1112.
- Mitchell RA & Berger AJ (1975). Neural regulation of respiration. *Am Rev Resp Dis* **111**, 206–224.
- Mithoefer JC (1965). Breath holding. In *Handbook of Physiology*, section 2, Eds. Fenn WO, Rahn H *Respiration*, pp. 1011–1025. American Physiological Society, New York.
- Nathan PW & Sears TA (1960). Effects of posterior root section on the activity of some muscles in man. *J Neurol Neurosurg Psychiat* **23**, 10–22.
- Noble MIM, Eisele JR, Frankel HL, Else W & Guz A (1971). The role of the diaphragm in the sensation of holding the breath. *Clin Sci* **41**, 275–283.
- Noble MIM, Eisele JH, Trenchard D & Guz A (1970). Effect of selective peripheral nerve blocks on respiratory sensations. In *Breathing: Hering–Breuer Centenary Symposium*, ed. Porter R, pp. 233–247. J & A Churchill (Longmans Group), London.
- Nunn JF (1987). *Applied Respiratory Physiology*. Butterworth Ltd, London.
- Orem J (1989). Behavioral inspiratory inhibition: inactivated and activated respiratory cells. *J Neurophysiol* **62**, 1069–1078.
- Orem J & Netick A (1986). Behavioral control of breathing in the cat. *Brain Res* **366**, 238–253.
- Passino C, Sleight P, Valle F, Spadacini G, Leuzzi S & Bernardi L (1997). Lack of peripheral modulation of cardiovascular central oscillatory autonomic activity during apnea in humans. *Am J Physiol* **272**, H123–H129.
- Paulev PE (1969). Respiratory and cardiovascular effects of breath holding. *Acta Physiol Scand supplement* **324**, 1–150.
- Pawelczyk JA & Levine BD (1995). Cardiovascular variability without ventilation: evidence for centrally mediated low frequency oscillations. *FASEB J* **9**, A840.
- Piepoli M, Sleight P, Leuzzi S, Valle F, Spadacini G, Passino C, Johnston J & Bernardi L (1997). Origin of respiratory sinus arrhythmia in conscious humans. *Circ J Am Heart Soc* **95**, 1813–1821.
- Porter R (1970). *Breathing: Hering–Breuer Centenary Symposium*. J. & A Churchill (Longmans Group), London.
- Rigg JR, Rebeck AS & Campbell EJM (1974). A study of factors influencing relief of discomfort in breath-holding in normal subjects. *Clin Sci Mol Med* **47**, 193–199.
- Road JD (1990). Phrenic afferents and ventilatory control. *Lung* **168**, 137–149.
- Rodbard S (1947). The effect of oxygen, altitude and exercise on breath-holding time. *Am J Physiol* **150**, 142–148.
- Schneider EC (1924). A comparison of three types of anoxemia. *Military Surgeon* March, 328–339.
- Schneider EC (1930). Observations on holding the breath. *Am J Physiol* **94**, 464–470.
- Schneider EC & Truesdell D (1923). The circulatory responses of man to a sudden and extreme anoxemia. *Am J Physiol* **65**, 379–385.
- Sears TA (1966). The respiratory motoneurone: integration at a spinal segmental level. In *Breathlessness*, ed. Howell JBL & Campbell EM, pp. 33–47. Blackwell Scientific Publications, Oxford.
- Shea SA, Andres LP, Shamon DC, Guz A & Banzett RB (1993). Respiratory sensations in subjects who lack a ventilatory response to CO₂. *Resp Physiol* **93**, 203–219.
- Shea SA (1996). Behavioural and arousal-related influences on breathing in humans. *Exp Physiol* **81**, 1–26.
- Sterba JA & Lundgren CE (1985). Diving bradycardia and breath-holding time in man. *Undersea Biomed Res* **12**, 139–150.
- Stevens CD, Ferris EB, Webb JP, Engel GL & Logan M (1946). Voluntary breath holding 1. Pulmonary gas exchange during breath holding. *J Clin Invest* **25**, 723–728.
- Supinski GS, Dick T, Stofan D & DiMarco AF (1993). Effects of intraphrenic injection of potassium on diaphragm activation. *J Appl Physiol* **74**, 1186–1194.

- Supinski GS, DiMarco AF, Hussein F & Altose MD (1989). Alterations in respiratory muscle activation in the ischemic fatigued canine diaphragm. *J Appl Physiol* **67**, 720–729.
- Tawadrous FD & Eldridge FL (1974). Posthyperventilation breathing patterns after active hyperventilation in man. *J Appl Physiol* **37**, 353–356.
- Trzebski A & Smietanowski M (1996). Cardiovascular periodicities in healthy humans in the absence of breathing and under reduced chemical drive of respiration. *J Auton Nerv Syst* **57**, 144–148.
- Trzebski A, Smietanowski M & Zebrowski J (2001). Repetitive apneas reduce nonlinear dynamical complexity of the human cardiovascular control system. *J Physiol Pharmacol* **52**, 3–19.
- Valentinuzzi ME & Geddes LA (1974). The central component of the respiratory heart-rate response. *Cardiovasc Res Cent Bull* **12**, 87–103.
- Ward SA, Macias D & Whipp BJ (2001). Is breath-hold time an objective index of exertional dyspnoea in humans? *Eur J Appl Physiol* **85**, 272–279.
- Wasserman K, Whipp BJ, Kayal SN & Cleary MG (1975). Effect of carotid body resection on ventilatory and acid base control during exercise. *J Appl Physiol* **39**, 354–358.
- Whitelaw WA, McBride B, Amar J & Corbet K (1981). Respiratory neuromuscular output during breath holding. *J Appl Physiol* **50**, 435–443.
- Whitelaw WA, McBride B & Ford GT (1987). Effect of lung volume on breath holding. *J Appl Physiol* **62**, 1962–1969.

Acknowledgements

I am very grateful to Dr Tom Clutton-Brock for his support and encouragement to Hannah Cooper and Jason Rutherford for their help and to John Ashworth, Mathew Bridge, Anna-Liese Acklam, Robert Dobson, Mark Scoble, Owen Kiely, Roy Jentjens, Katherine Horner, Rachel Felton, Kerry Pearn and Rachel Rickard for breath-holding.